

X-ray spectral diagnostics of activity in massive stars

David H. Cohen¹, Emma E. Wollman², & Maurice A. Leutenegger³

¹Department of Physics and Astronomy, Swarthmore College,
500 College Ave., Swarthmore, Pennsylvania, 19081, USA
email: cohen@astro.swarthmore.edu

²Department of Physics, California Institute of Technology,
Pasadena, California, 91125, USA
email: ewollman@caltech.edu

³NASA/Goddard Spaceflight Center,
Laboratory for High Energy Astrophysics,
Code 662, Greenbelt, Maryland, 20771, USA
email: maurice.a.leutenegger@nasa.gov

Abstract. X-rays are direct evidence of instabilities, time-variable structure, and shock heating in the winds of O stars. The observed broad X-ray emission lines provide information about the kinematics of shock-heated wind plasma and their shapes provide information about wind absorption, and thus about the wind mass-loss rates. Mass-loss rates determined from X-ray line profiles are not sensitive to density-squared clumping effects, and indicate mass-loss rate reductions of factors of 3 to 6 over traditional diagnostics that suffer from density-squared effects. Broad-band X-ray spectral energy distributions also provide mass-loss rate information via soft X-ray absorption signatures. In some cases, the degree of wind absorption is so high, that the hardening of the X-ray SED can be quite significant. We discuss these results as applied to the early O stars ζ Pup (O4 If), 9 Sgr (O4 V), and HD 93129A (O2 If*).

Keywords. line: formation, shock waves, stars: winds, x-rays: stars

1. Introduction

Activity in cool stars involves high-energy – usually magnetic – processes, X-rays, time variability, extended atmospheres and outflows. In normal O stars, the role of magnetic fields is more restricted and solar-type magnetic dynamos are generally thought not to operate. However, time-variable high-energy processes and their X-ray emission, associated with these stars’ powerful radiation-driven winds, are ubiquitous manifestations of activity in massive stars. Here we discuss the properties of the X-ray observations at high spectral resolution in three O and employ the observed X-rays as diagnostics of the activity in the outer atmospheres of these stars – specifically the shock heating and of their wind mass-loss rates.

We restrict the discussion of X-ray spectral diagnostics of activity to “normal” massive stars, where binarity and the associated colliding wind shock (CWS) X-ray emission and magnetically channeled wind shock (MCWS) X-ray emission is absent or negligible. The dominant paradigm for X-ray production in normal O and early-B stars is the embedded wind shock (EWS) scenario, where shock-heated regions are distributed throughout the stellar wind. The specific mechanism for EWSs is usually assumed to involve the line-driving instability (LDI), either in a self-excited mode (Owocki, Castor, & Rybicki 1988) or in a mode where the instability is seeded by perturbations at the base of the wind (Feldmeier, Puls, & Pauldrach 1997). At high spectral resolution, the X-ray emission from

the EWS scenario displays broad emission lines, from which the kinematics of the shock-heated plasma can be determined. Furthermore, these broad lines have a characteristic blue-shifted and asymmetric shape due to attenuation of the X-rays as they propagate through the bulk, cool wind component in which photoelectric absorption – a continuum process – preferentially attenuates red-shifted line photons emitted in the far hemisphere of the wind, while leaving blue-shifted line photons emitted from the near hemisphere much less attenuated. This line profile phenomenology allows us – via statistical fitting of empirical line profile models – to test the predictions of the LDI simulations via the kinematic information and also to measure the wind mass-loss rates, via the attenuation signatures of the profiles.

The morphology of high-resolution X-ray spectra of normal massive stars differs in some obvious ways from those of low-mass coronal sources, as we show in Fig. 1, where we compare the O supergiant ζ Pup to the coronal G star, Capella. The O star’s spectrum is harder, overall, than the G star’s, however this is due not to higher plasma temperatures, but rather to the effects of wind absorption, consistent with the X-rays arising in the dense stellar wind of the O star. This broadband view of the X-ray spectra also shows quite obviously that the emission lines in the O star are much broader than the (unresolved) lines in the G star, as the EWS scenario predicts. In the next section, we show how quantitative information can be derived from the Doppler broadened X-ray emission lines.

2. The X-ray line profile model applied to ζ Pup

To extract information from individual resolved line profiles, we fit a simple, empirical wind-shock model informed by the LDI simulations, in which numerous shock-heated regions are distributed throughout the wind above some shock onset radius, R_o , with local emission measure that we assume to scale with the local ambient wind density




Figure 1. *Chandra* medium energy grating (MEG) spectra of the O4 If star, ζ Pup (top) and for comparison, of the G star, Capella (bottom). The spectrum of the O star is harder (strongest lines at shortest wavelengths) but by comparing H-like and He-like line strengths (e.g. of Mg at 8.42 Å and 9.2 Å, respectively) it is evident that the higher temperature plasma is found on the G star. Finally, note that the emission lines are unresolved in the Capella spectrum and are significantly broadened in the ζ Pup spectrum.

squared (Owocki & Cohen 2001). The kinematic profile of the X-ray plasma is assumed to trace the same beta-velocity law that describes the bulk wind. This assumption is based on the results of numerical simulations of EWSs that show accelerated pre-shock wind streams being decelerated back down the local ambient wind velocity (Feldmeier, Puls, & Pauldrach 1997, Runacres & Owocki 2002). The attenuation due to continuum opacity in the bulk wind in which the shock-heated plasma is embedded is described by the characteristic optical depth parameter, $\tau_* \equiv \kappa \dot{M} / 4\pi R_* v_\infty$, where κ is the (wavelength dependent) opacity of the bulk wind, \dot{M} is the wind mass-loss rate, R_* is the stellar radius, and v_∞ is the wind terminal velocity.

For each emission line in the *Chandra* spectrum of an O star, we can fit this empirical profile model and derive best-fit values of R_o and τ_* by minimizing the C statistic, and place confidence limits on them via the $\Delta\chi^2$ formalism applied to the C statistic. For ζ Pup (summarizing the results published in Cohen et al. 2010), we find – for 16 lines and line complexes in the *Chandra* grating spectrum (three representative lines and their best-fit profile models are shown in Fig. 2) – a universal value for the shock-onset radius of $R_o \approx 1.5 R_*$, which is consistent with numerical simulations of the LDI. We also find a range of characteristic optical depths, τ_* , for the 16 emission lines, consistent with the expected wavelength trend in the atomic opacity. By calculating a detailed opacity model, and assuming standard values for the stellar radius and wind terminal velocity, we fit the ensemble of characteristic optical depths to find a best-fit mass-loss rate, via $\dot{M} = 4\pi R_* v_\infty \tau_*(\lambda) / \kappa(\lambda)$. The values of the onset radius, R_o , and of the characteristic optical depths, τ_* , are shown in Fig. 3. The panel with the τ_* values also shows the best-fit model of the wavelength-dependent optical depths, from which we derive a mass-loss rate of $3.5 \pm 0.3 \times 10^{-6} M_\odot \text{ yr}^{-1}$.

In closing this section, we emphasize that the mass-loss rate determination from the X-ray profiles represents a factor of roughly three reduction from the traditional H α -derived mass-loss rate for which clumping is ignored (Markova et al. 2004). And that this modest reduction in the mass-loss rate is consistent with newer determinations using H α and radio and IR free-free excesses with *do* account for clumping (Puls et al. 2006). Also, we note that for a high signal-to-noise *Chandra* spectrum with a lot of emission lines, like that of ζ Pup, the statistical error on the derived mass-loss rate is small (about 10%). But that the actual uncertainty is dominated by uncertainty in the wind opacity model, which in turn is dominated by uncertainty in the elemental abundances. Individual elemental abundances do not have a large effect, but the overall metallicity does. The model we




Figure 2. Fits to three lines in the *Chandra* spectrum of ζ Pup. From left to right: the Ly α line of Mg XII at 8.42 Å, the Fe XVII line at 15.01 Å, and the Ly α line of O VIII at 18.97 Å. The vertical dashed lines in each panel represent the laboratory rest wavelengths of each transition, while the flanking dotted lines represent the Doppler shifts associated with the wind terminal velocity. The characteristic broad, blue shifted, and asymmetric profile shapes are evident, as is an increase in the shift and asymmetry with wavelength, as is expected from the form of the continuum opacity of the bulk wind, which generally increases with wavelength. The characteristic optical depths of these three lines are roughly $\tau_* = 1, 2,$ and $3,$ respectively.

use in this paper (and which was used in Cohen et al. 2010) uses subsolar metallicity and C, N, and O abundances altered from the usual ratio by CNO processing. If future abundance determinations are made which supersede the current ones, the mass-loss rate should be rescaled in inverse proportion to the metallicity adjustment (more metals cause higher opacity which would then require lower wind column densities and so lower mass-loss rates).

3. Other O stars

We can apply the same type of line profile analysis to other O stars observed with the *Chandra* grating spectrometer. Here we present preliminary analysis of the early O main sequence star, 9 Sgr, at the center of the Lagoon Nebula, and the very early O supergiant, HD 93129A, in Tr 14 in Carina. Both stars have binary companions, but in neither case are the emission lines in the grating spectrum significantly contaminated by the harder X-rays associated with CWB X-rays.

There are seven lines and line complexes in the 9 Sgr *Chandra* grating spectrum with high enough signal-to-noise for line profile modeling to provide meaningful constraints. The mean shock onset radius is $R_o = 1.4 R_*$, consistent with the EWS scenario. The ensemble of τ_* values can be fit, given a model of the bulk wind opacity (which we calculate assuming solar abundances), to derive a mass-loss rate. We find a mass-loss rate of $\dot{M} = 4.2 \times 10^{-7} M_\odot \text{ yr}^{-1}$, which represents a factor of six reduction over the traditional mass-loss rate derived from H α fitting, assuming a smooth wind. In Fig. 4 we show the R_o and τ_* results.

The O2 If* star, HD 93129A, is the earliest O star and, according to Taresch et al. 1997, has the highest mass-loss rate of any O star, with $\dot{M} = 1.8 \times 10^{-5} M_\odot \text{ yr}^{-1}$. More recent modeling (though also ignoring clumping effects) gives $\dot{M} = 2.6 \times 10^{-5} M_\odot \text{ yr}^{-1}$ and a wind terminal velocity of $v_\infty = 3200 \text{ km s}^{-1}$ (Repolust, Puls, & Herrero 2004). This extremely strong and dense stellar wind provides an interesting test of the EWS scenario for X-ray production in O stars. Indeed, the *Chandra* spectrum is quite hard, which if interpreted as a temperature effect would make an EWS interpretation implausible. However, the H-like Si line strength is very weak, compared to the He-like




Figure 3. Results from fitting the wind-profile model to the emission lines in the *Chandra* spectrum of ζ Pup. From left to right: the shock onset radii, R_o , which are consistent with a universal value of $1.5 R_*$; the wind terminal velocities for the five strongest, unblended lines in the spectrum, which are consistent with the value for the bulk wind, of $v_\infty = 2250 \text{ km s}^{-1}$, derived from UV spectra (this value is represented by the horizontal line, while the cross hatched region is the 68% confidence limit on the mean value of the five fitted terminal velocities shown as points with error bars); and the τ_* values from each of the 16 fitted line profiles. This last panel shows that a constant value of the characteristic optical depth provides a poor fit, as does a model that incorporates the continuum opacity of the bulk wind but assumes a high value for the mass-loss rate ($8.3 \times 10^{-6} M_\odot \text{ yr}^{-1}$), while a model with the mass-loss rate as a free parameter provides a good fit, with a mass-loss rate of $3.5 \times 10^{-6} M_\odot \text{ yr}^{-1}$.

Si line strength, indicating a plasma emission temperature of no more than 8 million K, which is consistent with LDI simulations of wind shocks. The hardness of the X-ray spectrum appears instead to be due to severe attenuation of the soft X-ray emission by both the interstellar medium and the star’s own wind.

Because of the absent soft X-rays, there are only four lines and line complexes in the *Chandra* grating spectrum available for fitting. We show the strongest unblended single line, the Ly α line of Mg XII at 8.42 Å, in Fig. 5, along with the shock onset radii and the characteristic optical depths. The latter plot has the mass-loss rate fit superimposed on the τ_* values. For this star, too, we find a modest mass-loss rate reduction of roughly a factor of four over the value derived from H α fitting assuming no clumping.

4. Broadband X-ray properties

Given the strong wind absorption in the X-ray spectra of early O stars, we have modeled the broadband spectral energy distributions using simple one- and two-temperature thermal emission spectral models (e.g. APEC Smith et al. 2001) along with wind attenuation, using the newly published radiation transport model, *windtabs* (Leutenegger et al. 2010). This model accounts for the spatial distribution of the emitting plasma within the absorbing wind, and thus has a much more gradual decrease of transmission vs. fiducial optical depth than is seen in the exponential absorption model that describes an absorbing medium in between the background emitter and the observer, as is employed in interstellar absorption models. The *windtabs* absorption model also employs a realistic photoionization opacity model that includes partially ionized metals and fully ionized H and He. We fit the *Chandra* zeroth-order spectrum (a CCD low-resolution spectrum) of HD 93129A with this APEC and *windtabs* model and find a low plasma temperature of 0.6 keV – fully consistent with the LDI simulation results – and a significant wind column density, corresponding to a mass-loss rate of $\dot{M} = 8 \times 10^{-6} M_{\odot} \text{ yr}^{-1}$, which is consistent with the value we find from fitting the individual line profiles (shown in the third panel of Fig. 5). We show the HD 93129A zeroth-order spectrum and best-fit APEC and *windtabs* model in Fig. 6, along with the grating spectrum of the star, which shows the low H-like/He-like line ratios, indicative of low plasma temperatures.




Figure 4. Results from fitting the wind-profile model to the emission lines in the *Chandra* spectrum of 9 Sgr. The shock onset radii (left) are consistent with a value of $R_o = 1.4 R_{\odot}$, while the characteristic optical depths (right) are well fit by a model that has a mass-loss rate of $\dot{M} = 4.2 \times 10^{-7} M_{\odot} \text{ yr}^{-1}$ (solid line), a factor of six below the unclumped H α mass-loss rate (dotted line).

5. Conclusions

Massive stars with their powerful, dense, radiation-driven winds show a significant amount of activity: time-variable structure, dissipation of energy in their extended atmospheres, and X-ray emission. The X-ray emission, when analyzed in detail, shows that the properties of the hot plasma that produces the emission are consistent with the embedded wind shock scenario. Specifically, the hot plasma shares the same kinematic profile as the bulk wind (with terminal velocities in excess of 2000 km s^{-1}), that it is spatially distributed throughout the wind above an onset radius of roughly $R_o = 1.5 R_*$, and – from broadband modeling – that the plasma temperatures are less than 10 million K, in accord with the predictions of LDI simulations of EWSs. These relatively low temperatures can be reconciled with the relatively hard observed spectra by taking wind attenuation of the soft X-rays into account. When we model the effect of wind attenuation on individual emission lines, we find that their modestly blue-shifted and asymmetric profiles can be reproduced using mass-loss rates that are lower by a factor of 3 to 6 compared to traditional mass-loss rates that ignore clumping (and are consistent with newer determinations that account for the clumping). And furthermore, we find that when we model the broadband spectral properties and account for the effects of wind attenuation using a realistic radiation transport model in conjunction with a realistic opacity model, we derive similar mass-loss rate values.

Finally, we note that in this short paper we do not have the space to discuss in detail the possible role of *porosity* in generating the only modestly blue-shifted and asymmetric profiles. Porosity arises from clumping on very large scales, where individual clumps are optically thick to X-ray photoelectric absorption. Our modeling suggests that porosity does not need to be invoked in order to explain the observed X-ray properties of the early O stars we discuss here. Their properties are well explained by modest mass-loss rate reductions. Furthermore, porosity requires clumping, by definition (but not the other way around). So, once clumping is invoked, and the density-squared diagnostics are adjusted accordingly, there is no longer any need to invoke porosity to explain the data. This and other aspects of porosity are addressed in the end-of-session discussion, later in these proceedings.

References

- Cohen, D.H., Leutenegger, M.A., Wollman, E.E., Zsargoó, J., Hillier, D.J., Townsend, R.H.D., Owocki, S.P. 2010, *MNRAS*, 405, 2391
 Feldmeier, A., Puls, J., Pauldrach, A.W.A. 1997, *A&A*, 322, 878
 Leutenegger, M.A., Cohen, D.H., Zsargoó, J., Martell, E.M., MacArthur, J.P., Owocki, S.P., Gagné, M., Hillier, D.J. 2010, *ApJ*, 719, 1767




Figure 5. Results from fitting the wind-profile model to the emission lines in the *Chandra* spectrum of HD 93129A: The Mg XII Ly α line (left) along with the R_o values (center) and the τ_* values (right) along with the best-fit mass-loss rate model (solid line) and for comparison, the traditional unclumped H α mass-loss rate.

- Markova, N., Puls, J., Repolust, T., Markov, H. 2004, *A&A*, 413, 693
 Owocki, S.P., Castor, J.I., Rybicki, G.B. 1988, *ApJ*, 335, 914
 Owocki, S.P., Cohen, D.H. 2001, *ApJ*, 559, 1108
 Puls, J., Markova, N., Scuderi, S., Stanghellini, C., Taranova, O.G., Burnley, A.W., Howarth, I.D. 2006, *A&A*, 454, 625
 Repolust, T., Puls, J., Herrero, A. 2004, *A&A*, 415, 349
 Runacres, M.C., Owocki, S.P. 2002, *A&A*, 381, 1015
 Smith, R.K., Brickhouse, N.S., Liedahl, D.A., Raymond, J.C. 2001, *ApJ*, 556, L91
 Taresch, G., et al. 1997, *A&A*, 321, 531




Figure 6. The zeroth-order, low-resolution *Chandra* spectrum of HD 93129A, with the best-fit thermal emission model attenuated by stellar wind absorption. This modeling shows that the emission temperature of the plasma is relatively low – $kT = 0.6$ keV – while the affects of wind attenuation are significant; explaining the observed hardness. The grating spectrum of this star is shown in the right-hand panel. Note that while the spectrum is quite hard (no significant emission longward of 10 \AA), the Si xiv Ly α line at 6.18 \AA is very weak compared to the Si XIII complex near 6.7 \AA . This very low H-like/He-like line ratio is indicative of plasma with a temperature of no more than 8 million K (0.7 keV).